

The success of the Gobiidae in tropical Pacific insular streams

P. A. RYAN*

Biology Department
School of Pure and Applied Sciences
University of the South Pacific
P.O. Box 1168, Suva, Fiji

*Present address: c/- West Coast Regional Council,
P.O. Box 66, Greymouth, New Zealand

Abstract The high islands of the tropical Pacific possess a sparse native fish fauna dominated by the family Gobiidae. Evidence of this domination is presented, together with possible reasons for the success of the Gobiidae. Factors which may have contributed include a marine larval stage, euryhalinity, small size, an excellent climbing ability, a range of trophic level from carnivory through omnivory to herbivory, frequent lack of a gas bladder, and an associated bottom-living life style. The evolution of streams in the high islands of the tropical Pacific is relatively recent, thus there has been little time for colonisation to occur. Furthermore, estuaries in many islands are poorly developed, thereby placing euryhaline estuarine species at a competitive disadvantage.

Keywords Gobiidae; evolution; freshwater; distribution; colonisation; Pacific; islands; reproductive strategies; euryhalinity

INTRODUCTION

Native primary and secondary freshwater fish are absent from the tropical Pacific (Herre 1940; Myers 1953; Berra 1981) and although the faunal biomass

of insular inland waters may be similar to those of comparable continental waters, the species diversity is much lower (Maciolek 1984). Of the larger islands in the tropical Pacific, Fiji has been one of the best studied. The group was visited by Wilkes and the United States Exploring Expedition of 1838–1842, by the “Challenger” Expedition in 1874 and by Gardiner (in Boulenger 1897).

Publications on fish collections from the group have been made by Nutting (1924), Whitley (1927), Fowler (1928, 1929, 1959), and Herre (1936). More recently there have been visits from the Royal Ontario Museum (1983) and the Smithsonian Institution (1982, 1985). In the latter visits most of the sampling effort was directed at marine habitats. Ryan (1980)* published a checklist of Fiji’s brackish and freshwater fish and some new records (1981*). Information is also available from the Samoas (Jordan & Seale 1906; Schultz 1943; Wass 1984)*, Vanuatu (Baker 1929*; Ryan 1986*; and author’s collections), Palau (Bright & June 1981*), the Cook Islands (author’s observations and Jellyman in press*). For comparison with a continental landmass, information on a collection of fishes by Allen & Hoese (1980*), from the Jardine River, Cape York Peninsula, Australia, is also included. Unfortunately, no checklists could be obtained from north east Queensland which would have provided a more meaningful comparison with the tropical Pacific. Information from asterisked authors is summarised in Table 1. Many workers (following Springer 1983; Hoese 1984) split gobioid fishes into the Eleotridae and the Gobiidae but for the purposes of this paper I have followed Miller (1973) as the arguments apply equally well to both eleotrids and gobiids. For other families I have followed the classification used by Masuda et al. (1984).

RESULTS

The carnivorous sicydiine *Sicyopus zosterophorum*, not previously reported from Fiji, was collected by the author on Viulevu, Vanualevu, and Taveuni. The presence of the ophichthid eel, *Achirophichthys*

Table 1 Families and numbers of species represented in some selected tropical insular streams. Cape York, Australia, is included for comparison. Percentage representation of each family is given in brackets.

	Fiji	Vanuatu	Samoa	Cook Islands	Palau	Cape York
CLASS Chondrichthyes						
Order Lamniformes						
Family Carcharhinidae	1 (1.3)	--	--	--	--	--
Order Rajiformes						
Family Dasyatidae	1 (1.3)	--	--	--	--	--
CLASS Osteichthyes						
Order Elopiformes						
Family Elopidae	2 (2.5)	--	2 (4.1)	--	1 (2.5)	1 (3.3)
Family Albulidae	1 (1.3)	--	1 (2)	--	1 (2.5)	--
Order Anguilliformes						
Family Anguillidae	4 (5.0)	3 (10.3)	3 (6.1)	3 (30)	2 (5.0)	1 (3.3)
Family Moringuidae	1 (1.3)	--	1 (2)	--	1 (2.5)	--
Family Muraenidae	1 (1.3)	--	1 (2)	--	1 (2.5)	--
Family Muraenesocidae	1 (1.3)	--	--	--	--	--
Family Ophichthidae	3 (3.8)	1 (3.4)	1 (2)	--	--	--
Order Osteoglossiformes						
Family Osteoglossidae	--	--	--	--	--	1 (3.3)
Order Gonorhynchiformes						
Family Chanidae	1 (1.3)	--	1 (2)	1 (10)	--	--
Order Siluriformes						
Family Ariidae	--	--	--	--	--	1 (3.3)
Family Plotosidae	2 (2.5)	--	1 (2)	--	--	4 (13.3)
Order Beloniformes						
Family Belonidae	--	--	--	--	--	1 (3.3)
Family Hemiramphidae	1 (1.3)	--	1 (2)	--	--	--
Order Syngnathiformes						
Family Syngnathidae	6 (7.5)	2 (6.9)	4 (8.2)	--	4 (10)	--
Order Lophiiformes						
Family Antennariidae	--	--	--	--	1 (2.5)	--
Order Perciformes						
Family Atherinidae	--	--	--	--	--	1 (3.3)
Family Melanotaeniidae	--	--	--	--	--	6 (20)
Family Mugilidae	3 (3.8)	3 (10.3)	1 (2)	1 (10)	1 (2.5)	?
Family Centropomidae	--	--	--	--	--	1 (3.3)
Family Ambassidae	1 (1.3)	1 (3.4)	1 (2)	--	1 (2.5)	3 (10)
Family Serranidae	1 (1.3)	--	--	--	--	--
Family Kuhliidae	4 (5.0)	3 (10.3)	2 (4.1)	?	2 (5.0)	--
Family Apogonidae	1 (1.3)	--	1 (2)	--	--	1 (3.3)
Family Carangidae	1 (1.3)	1 (3.4)	2 (4.1)	--	--	--
Family Leiognathidae	1 (1.3)	--	--	--	--	--
Family Gerreidae	2 (2.5)	1 (3.4)	3 (6.10)	--	--	--
Family Monodactylidae	1 (1.3)	--	1 (2)	--	--	--
Family Lutjanidae	1 (1.3)	1 (3.4)	1 (2)	--	--	?
Family Teraponidae	2 (2.5)	1 (3.4)	1 (2)	--	--	2 (6.6)
Family Scatophagidae	1 (1.3)	1 (3.4)	--	--	1 (2.5)	--
Family Toxotidae	--	--	--	--	--	1 (3.3)
Family Siganidae	1 (1.3)	--	--	--	--	--
Family Gobiidae	33 (41.3)	10 (34.5)	18 (36.7)	5 (50)	23 (57.5)	5 (16.6)
Order Tetraodontiformes						
Family Tetraodontidae	2 (2.5)	--	2 (4.1)	--	--	--
Order Synbranchiformes						
Family Synbranchidae	--	--	--	--	1 (2.5)	1 (3.3)
Total species	80	29	49	10	40	30

kamperi, predicted to be found in Fiji (Ryan 1980), was confirmed from electro-fished specimens caught in Vitilevu by the Fisheries Department during 1987 and identified by the author. Dawson (1984), added the syngnathids *Microphis brevidorsalis*, *Microphis argulus*, and *Microphis brachyurus* to the Fijian, Samoan, and Vanuatuan fauna, respectively. These additions have been incorporated in the families represented in Tables 1 and 2. A checklist of the freshwater fish of the Cook Islands is given in Table 3.

DISCUSSION

Despite Fiji's distance from centres of origin (probably the Philippines), Fiji has a substantial fresh and brackish water fish fauna of at least 80 species, represented by 28 families (Ryan 1980). This diversity is much greater than the 30 species reported from the Cape York Peninsula, Australia, by Allen & Hoese (1980), the 40 species from Palau (Bright & June 1981), or the 29 species reported from Vanuatu (Baker 1929; this paper). This may just indicate the intensity with which Fiji has been studied or could be a reflection of the nearly four degrees of latitude covered by the group which has enhanced the chances of marine tolerant, or freshwater species with marine larval stages, reaching the islands. Vitilevu is at least 40×10^6 years old (Kroenke 1984) which has given

ample time for both colonisation and speciation. Despite this, with the one obvious exception of the family Gobiidae, little speciation has occurred. Gobiids account for 41% of the native Fijian stream fish fauna and fill a wide variety of niches. There are at least 4 endemics out of the 33 brackish and freshwater species and a number still remain undescribed. It is likely that there are 10 or more endemic gobies accounting for around 30% of the total gobiid fauna and around 12.5% of the total brackish/freshwater fauna.

Information from other tropical Pacific islands is sparse. In Palau over half of the freshwater fish fauna is composed of gobiids although there are only two known endemic species (Bright & June 1981). Of the 29 species known from Vanuatu, 10 (35%) are gobiids. At least one gobiid is endemic (Ryan 1986). Samoan gobiids make up 37% of the freshwater fishes (18 out of 49 species). In the Cook Islands, gobies make up 50% of a depauperate stream fauna (including one yet to be described species of *Stiphodon* in author's collection). In Cape York, the gobiid component of the fauna represents only 16% of the total species, a contribution less than half that of any of the island groups considered.

The question arises as to why there has not been more speciation amongst the other families on oceanic islands? (*Mesopristes kneri*, an endemic Fijian

Table 2 Brackish and freshwater fish species reported from Vanuatu. "*" denotes observed or collected by the author; "+" indicates information from Baker (1929).¹ Dawson (1984).

Species	Family Carangidae
Family Anguillidae	<i>Caranx sexfasciatus</i> +
<i>Anguilla megastoma</i> +	Family Gerreidae
<i>Anguilla marmorata</i> +	<i>Gerres filamentosus</i> +
<i>Anguilla obscura</i> +	Family Lutjanidae
Family Ophichthidae	<i>Lutjanus argentimaculatus</i> +
<i>Ophichthus polyophthalmus</i> +	Family Teraponidae
Family Syngnathidae	<i>Mesopristes argenteus</i> +
<i>Microphis leiaspis</i> *	Family Scatophagidae
<i>Microphis retzii</i> *	<i>Scatophagus argus</i> +
<i>Microphis brachyurus</i> ¹	Family Gobiidae
Family Mugilidae	<i>Guavina gyrinoides</i> +
<i>Liza macrolepis</i> +	<i>Ophiocara porocephala</i> +
<i>Crenimugil crenilabis</i> +	<i>Ophiocara macrolepidota</i> +
<i>Cestraeus plicatilis</i> +	<i>Ophioeleotris aporos</i> +
Family Ambassisidae	<i>Gobius sp.</i> +
<i>Ambassis urotaenia</i> +	<i>Awaous ocellaris</i> +
Family Kuhlidae	<i>Stenogobius genivittatus</i> +
<i>Kuhlia repustris</i> +	<i>Sicyopterus micrurus</i> +
<i>Kuhlia bilunulata</i> +	<i>Stiphodon "elegans"</i> *
<i>Kuhlia marginata</i> +	<i>Stiphodon astilbos</i> *

teraponid, is the only one of which I am aware). A reasonable hypothesis is that non-gobiids have been outcompeted by the gobies to the extent that speciation has not been able to take place. This competition takes place on a variety of fronts. In many gobiids the adults breed in freshwater, only the eggs or fry making the journey to the sea (Miller 1984). This passive transport requires little energy expenditure. By contrast, most, if not all, of the other families encountered in tropical Pacific insular streams (TPIS) require the return of reproductively mature adults to the sea or estuaries for breeding to occur. A number of metabolic changes take place before the migration and as a result of this journey, less energy is available for reproduction (Miller 1984). In those long-lived species which breed each year, energy must also be expended in the return upstream. Probable examples of this type of strategy would include the various *Kuhlia* species (Lewis & Hogan 1987). In energy terms, goby reproductive strategies may be more efficient than their competitors.

In Fijian freshwater gobies, most would appear to be *r* selected in marked contrast to the many *k* selected marine species. Female gobiids of the genera *Stiphodon* and *Sicyopterus* produce large numbers of tiny eggs and Fijians harvest the upriver migration of young larvae. Elsewhere, Maciolk (in McDowall 1988) considers that the sicydiine genera *Lentipes*, *Sicydium*, *Sicyopterus*, *Sicyopus*, and *Stiphodon* all lay eggs in freshwater with the young being swept to the sea and migrating back up rivers as larvae. I have collected a small upstream migrating school of transparent gobiid larvae in the lower reaches of Fiji's Rewa river (although it is not known where the adults spawn.) Tank reared, they were identified as *Hypseleotris guntheri*.

McDowall (1988) summarises our current knowledge of the breeding biology of a number of freshwater gobiids. Amphidromy (Myers 1949) seems to be the rule, rather than the exception, amongst gobiids inhabiting TPIS.

Species diversity increases in estuaries and lagoons and with this increase in diversity there are more predators, thus increasing the hazards of a catadromous life style. There is a relative lack of midwater predators in TPIS (author's unpublished data), so larvae drifting in streams face little predation although they are exposed to a full gamut of predators upon arrival in the estuary. At this point they are no more at risk than those *r* selected species which breed in estuaries (although they may be smaller). However, the adult breeding stock has not been exposed to the additional risks faced by downstream migration.

The largest gobiid encountered in TPIS is *Ophioeleotris aporos* which can reach 30 cm. Most gobiids are considerably smaller than this. Small size confers the ability to penetrate streamlets and utilise niches unavailable to marine families with relatively larger euryhaline representatives such as the Carangidae, Lutjanidae, or Serranidae. Smallness is also an advantage if food is in short supply, which would be the situation in depauperate streams on newly-formed islands. This does not explain why small specimens of predominantly marine species cannot compete for niche space with the TPIS Gobiidae. Estuarine species are usually capable of short forays into freshwater. When they do move upstream, large size confers an osmotic advantage over smaller individuals. The effect of size on oxygen demand can be described by the power function: $y = ax^b$ where y = oxygen uptake x = body weight

The exponent b for most fish has an average value of 0.86 (Brett & Groves 1979). Most gas exchange is through the gills (Hughes 1980) and so is ionic regulation (Evans 1980). The implication is that larger fishes require proportionately less oxygen than small fish. Hence, the branchial epithelium and rate of ionic exchange should also be proportionately smaller. Large grouper (*Epinephelus* spp.) are frequently found in the lower reaches of freshwater streams (author's observations and P. Kailola, pers. comm.) but are too big to find an adequate food supply or physically to enter many streams. Small specimens, which seem ideally suited to such habitats, do not occur there, perhaps because of the osmotic stress that they would encounter.

Another possible reason for the lack of speciation amongst the non-gobiid component of TPIS stems from the difference between oceanic islands and continental islands and continents. Most Pacific islands are geologically young. The Fijian island of Taveuni (< 1×10^6 years old) provides a graphic example of what many of these islands may have been like. On the east coast of Taveuni there is an almost constant slope of 15° stemming from a 1200 m mountain chain. There are few estuaries. In most instances, the water changes from fully fresh to fully salt in the matter of a few metres. Under these conditions, species that breed in freshwater and have a marine larval stage are favoured, as are those such as the Anguillidae which actually breed in the sea. Anguillids are the second most successful colonisers of TPIS (in terms of species diversity). Estuarine breeders will be at a great, if not impossible disadvantage. Weathering will eventually cause estuaries to form, but by then the majority of niches in freshwater will have already been colonised.

Streams on the east coast of Taveuni are characterised by large numbers of waterfalls. Gobies, because their pelvic fins are modified as a sucking disc, are at an advantage in colonising headwaters. Even young eleotridines, without a suctorial disc may still use their pelvic fins to aid in climbing. I have seen the gobiine *Parioglossus* sp. (which lacks a suctorial disc) climbing the glass walls of aquaria in this manner. Young streams often lack gravel and sand substrates simply because there has not been enough weathering to form them. This limits food supplies and reduces relief from the current. On Taveuni, with its 15° slope, fish without low energy anchoring devices must swim the entire time just to keep station. Again, this places them at a competitive disadvantage in comparison with the gobiids.

Most, if not all, TPIS gobiids are negatively buoyant. After collection with rotenone or quinaldine, few species float to the surface. Most gobies also lack a gas bladder, in marked contrast to the majority of non-gobiid estuarine species. The end result is that even in the absence of a suctorial disc, gobiids expend less energy keeping station than gas bladder equipped species.

Another advantage possessed by the Gobiidae is trophic. They cover a full spectrum of feeding strategies from herbivory through omnivory to carnivory. Many TPIS gobiids are herbivores or omnivores. In the Cook Islands for instance, three of the five gobiids recorded by the author (Table 3) were herbivorous. The other two were carnivores

with omnivorous tendencies (*Eleotris fuscus* and *Awaous ocellaris*). With such a suite of choices resident within the family, it is hardly surprising that some species are able to colonise the virgin habitat a recently formed stream offers. Once present, selection ensures that the resident gobies become better adapted to the environment.

The three herbivorous gobies (*Stiphodon "elegans"*, *Stiphodon "stevensoni"*, and *Sicyopterus "micrurus"*) are sympatric algal grazers and may reach very high densities in some streams. It is not known how they partition the algal resource. The same pattern of two *Stiphodon* and sometimes two *Sicyopterus* spp. is repeated in many other TPIS. In French Polynesia and on Vanualevu in Fiji I have observed interactions between four algal-grazing sicydiine species. The situation is made even more complicated by competition from algal-grazing molluscs!

The ideas presented above require further development, particularly the comments on goby reproductive strategies. However there is little doubt that the gobies are TPIS colonisers par excellence. Once the occupation has occurred it may be difficult for them to be displaced.

Genetic isolation between different stream populations and the original colonisers will favour speciation. The relative importance of the different contributory factors is unknown, but this could be a fruitful field of study for evolutionary biologists.

Table 3 The trophic status of fresh and brackish water species reported from the Cook Islands. "*" denotes collected by author, "+" denotes Jellyman pers. comm.

Species	Trophic status
Family Anguillidae	
<i>Anguilla megastoma</i> +	carnivore
<i>Anguilla marmorata</i> +	carnivore
<i>Anguilla obscura</i> +	carnivore
Family Chanidae	
<i>Chanos chanos</i> + ¹	herbivore
Family Mugilidae	
<i>Mugil cephalus</i> *	herbivore
Family Gobiidae	
<i>Awaous ocellaris</i> *	carnivore
<i>Stiphodon "elegans"</i> *	herbivore
<i>Stiphodon "stevensoni"</i> *	herbivore
<i>Sicyopterus "micrurus"</i> *	herbivore
<i>Eleotris fuscus</i> *	carnivore

¹Farmed in Lake Terotonue, Mitiaro. Juveniles or adults may ascend a short distance up some streams on Rarotonga.

ACKNOWLEDGMENTS

This paper is based on one presented to the third Indo-Pacific fish conference in Wellington. Thanks to Rick Winterbottom, Doug Hoes, Peter Miller, Mike Winterbottom, Helen Larson, Terry Donaldson, Bruce Collette, and Chris Paulin for helping me to refine these ideas over a beer or three. Special thanks also to John Maciolek and Bill Kenchington for reading the manuscript and Chris Paulin for insisting that I include Palau so as to avoid an acronym for Southern Hemisphere Insular Tropical Streams. The anonymous referees made some useful suggestions, most of which have been incorporated in the paper.

REFERENCES

- Allen, G. R.; Hoes, D. F. 1980: A collection of fishes from the Jardine River, Cape York Peninsula, Australia. *Journal of the Royal Society of Western Australia* 63 (2): 53-61.
- Baker, J. R. 1929: Man and animals in the New Hebrides. London, George Routledge & Sons Ltd. 200 p.

- Berra, T. M. 1981: An atlas of the freshwater fish families of the world. Lincoln and London. University of Nebraska Press. 197 p.
- Boulenger, G. A. 1897: A list of fishes obtained by Mr J. Stanley Gardiner at Rotuma, South Pacific Ocean. *Annual magazine of natural history, London* 6 (20): 371-374.
- Brett, J. R.; Groves, T. D. D. 1979: Physiological energetics. In: Hoar, W. S.; Randall, D. J.; Brett J. R. ed., Fish physiology vol. VIII. N. Y., Academic Press.
- Bright, G. R.; June, J. A. 1981: Freshwater fishes of Palau, Caroline. *Micronesica* 17: 107-111.
- Dawson, C. E. 1984: Revision of the genus *Microphis* Kaup (Pisces: Syngnathidae). *Bulletin of marine science* 35(2): 117-181.
- Evans, D. H. 1980: Osmotic and ionic regulation by freshwater and marine fishes. In: Ali, M. A. ed., Environmental physiology of fishes. New York and London, Plenum Press. 723 p.
- Fowler, H. W. 1928: The fishes of Oceania. *Memoirs of the Bernice P. Bishop Museum* 10. 540 p.
- 1929: Fishes obtained at Fiji in 1929. *Bernice P. Bishop Museum occasional papers* 9 (20): 113.
- 1959: Fishes of Fiji. Suva, Government of Fiji. 670 p.
- Herre, W. C. T. 1936: Fishes of the Crane Pacific Expedition. *Field Museum of Natural History, zoological series* 21: 14-72.
- 1940: Distribution of freshwater fishes in the Indo-Pacific. *Scientific monthly* 51: 165-168.
- Hoese, D. F. 1984: Gobioidae: Relationships. *American Society of Ichthyologists and Herpetologists, special publication number* 1: 760 p.
- Hughes, G. M. 1980: Morphometry of fish gas exchange organs in relation to their respiratory function. In: Ali, M. A. ed., Environmental physiology of fishes. New York and London, Plenum Press. 723 p.
- Jellyman, D. J. 1991: The biology of the shortfinned eel *Anguilla obscura* in Lake Terotonuc, Mitiaro, Cook Islands. *Pacific Science*. In press.
- Jordan, D. S.; Seale, A. 1906: The fishes of Samoa. Descriptions of the species found in the archipelago, with a provisional checklist of the fishes of Oceania. *Bulletin of the Bureau of Fisheries* 25: 173-455.
- Kroenke, L. 1984: Cenozoic tectonic development of the southwest Pacific. Commission for coordination of joint prospecting for mineral resources in South Pacific offshore areas. *South Pacific Applied Geoscience Commission technical bulletin* 6.
- Lewis, A. D.; Hogan, A. E. 1987: The enigmatic jungle perch recent research provides some answers. *South Pacific Commission newsletter* 40: 24-31.
- McDowall, R. M. 1988: Diadromy in fishes. Portland, Timber Press. 307 p.
- Maciolek, J. A. 1984: Exotic fishes in Hawaii and other islands of Oceania. In: Courtenay, W. R. (Jnr); Stauffer, J. R. (Jnr) ed., Distribution, biology and management of exotic fishes. Baltimore, Johns Hopkins University Press.
- Masuda, H.; Amaoka, K.; Araga, C.; Uyeno, T.; Yoshino, T. (eds.) 1984: The fishes of the Japanese Archipelago. Tokyo, Tokai University Press. Text volume. 437 p.
- Miller, P. J. 1973: The osteology and adaptive features of *Rhyacichthys aspro* (Telostei: Gobioidae) and the classification of gobioid fishes. *Journal of zoology* 171: 397-434.
- 1984: The Tokology of Gobioid Fishes. In: Potts, G. W.; Woolton, C. J. ed., Fish reproduction, strategies and tactics. London, Academic Press. 416 p.
- Myers, G. S. 1949: Salt tolerance of freshwater fish groups in relation to zoogeographical problems. *Bijdragen tot de Dierkunde* 28: 315-322.
- 1953: Paleogeographical significance of freshwater fish distribution in the Pacific. *Proceedings of the 7th Pacific Science Congress* 4: 38-48.
- Nutting, C. C. 1924: The fishes in the Fiji-New Zealand Expedition. *University of Iowa studies in natural history* 10 (5): 56-67.
- Ryan, P. A. 1980: A checklist of the brackish and freshwater fish of Fiji. *South Pacific journal of natural sciences* 1: 58-73.
- 1981: Records of three new freshwater fishes from the Fiji Islands. *Pacific science* 35 (1): 93-95.
- 1986: A new species of *Stiphodon* (Family Gobiidae: Sicydiaphiinae) from Vanuatu. *Proceedings of the second international conference on Indo-Pacific fishes*. 655-662.
- Schultz, L. P. 1943: Fishes of the Phoenix and Samoan Islands collected in 1939 during the expedition of the U. S. S. "Bushnell". *Bulletin of the United States National Museum* 180: 13-16.
- Springer, V. G. 1983: *Tyson belos*, new genus and species of western Pacific fish (Gobiidae, Xenisthminae), with discussion of gobioid osteology and classification. *Smithsonian contributions to zoology* 390: 140.
- Wass, R. C. 1984: An annotated checklist of the Fishes of Samoa. *National oceanic and atmospheric administration technical report. National Marine Fisheries Service*. SSRF781. 43 p.
- Whitley, G. P. 1927: A checklist of fishes recorded from Fijian waters. *Pan-Pacific Research Institute journal* 2 (1): 38.